

DEGENERATIONS OF GRADED COHEN-MACAULAY MODULES

NAOYA HIRAMATSU

ABSTRACT. We introduce a notion of degenerations of graded modules. In relation to it, we also introduce several partial orders as graded analogies of the hom order, the degeneration order and the extension order. We prove that these orders are identical on the graded Cohen-Macaulay modules if a graded ring is of graded finite representation type and representation directed.

1. INTRODUCTION

The notion of degenerations of modules appears in geometric methods of representation theory of finite dimensional algebras. In [15] Yoshino gives a scheme-theoretical definition of degenerations, so that it can be considered for modules over a Noetherian algebra which is not necessary finite dimensional. Now, a theory of degenerations is considered for derived module categories [10] or stable module categories [16]. The degeneration problem of modules has been studied by many authors [6, 12, 14, 15, 18, 19]. For the study, several order relations for modules, such as the hom order, the degeneration order and the extension order, were introduced, and the connection among them has been studied. In the previous paper [8], the author give the complete description of degenerations over a ring of even-dimensional simple hypersurface singularity of type (A_n) .

In the present paper we consider degenerations of graded Cohen-Macaulay modules over a graded Gorenstein ring with a graded isolated singularity. First we consider an order relation on a category of graded Cohen-Macaulay modules which is called the hom order (Definition 2.2). We shall show that it is actually a partial order if a graded ring is Gorenstein with a graded isolated singularity.

In section 3 we propose a definition of degenerations for graded modules and we state several properties on it. We show that if the graded ring is of graded finite representation type and representation directed, then the hom order, the degeneration order and the extension order are identical on the graded Cohen-Macaulay modules. We also consider stable analogue of degenerations of graded Cohen-Macaulay modules in section 4.

Date: February 8, 2013.

2000 Mathematics Subject Classification. Primary 16W50 ; Secondary 13D10.

Key words and phrases. degeneration, graded Cohen-Macaulay module, finite representation type.

2. HOM ORDER ON GRADED MODULES

Throughout the paper $R = \bigoplus_{i=0}^{\infty} R_i$ be a commutative Noetherian \mathbb{N} -graded Cohen-Macaulay ring with $R_0 = k$ a field of characteristic zero. A graded ring is said to be * local if the set of graded proper ideals has a unique maximal element. Thus R is * local since $\mathfrak{m} = \bigoplus_{i>0} R_i$ is a unique maximal ideal of R . We denote by $\text{mod}^{\mathbb{Z}}(R)$ the category of finitely generated \mathbb{Z} -graded modules whose morphisms are homogenous morphisms that preserve degrees. For $i \in \mathbb{Z}$, $M(i) \in \text{mod}^{\mathbb{Z}}(R)$ is defined by $M(i)_n = M_{n+i}$. Then $\text{Hom}_R(M, N(i))$ consisting of homogenous morphisms of degree i , and we set

$$^*\text{Hom}_R(M, N) = \bigoplus_{i \in \mathbb{Z}} \text{Hom}_R(M, N(i)).$$

For a graded prime ideal \mathfrak{p} of R , we denote by $R_{(\mathfrak{p})}$ a homogenous localization of R by \mathfrak{p} . For $M \in \text{mod}^{\mathbb{Z}}(R)$, take a graded free resolution

$$\cdots \rightarrow F_l \xrightarrow{d_l} F_{l-1} \rightarrow \cdots \rightarrow F_1 \xrightarrow{d_1} F_0 \rightarrow M \rightarrow 0.$$

We define an l th syzygy module $\Omega^l M$ of M by $\text{Im } d_l$. We say that a graded R -module M is said to be a graded Cohen-Macaulay R -module if

$$^*\text{Ext}_R^i(R/\mathfrak{m}, M) = 0 \quad \text{for any } i < d = \dim R.$$

In particular, this condition is equivalent to

$$^*\text{Ext}_R^i(M, \omega_R) = 0 \quad \text{for any } i > 0,$$

where ω_R is a * canonical module of R . We denote by $\text{CM}^{\mathbb{Z}}(R)$ the full subcategory of $\text{mod}^{\mathbb{Z}}(R)$ consisting of graded Cohen-Macaulay R -modules. By our assumption on R , $\text{mod}^{\mathbb{Z}}(R)$ and $\text{CM}^{\mathbb{Z}}(R)$ are Krull-Schmidt, namely each object can be decomposed into indecomposable objects up to isomorphism uniquely. For $M \in \text{mod}^{\mathbb{Z}}(R)$ we denote by $h(M)$ a sequence $(\dim_k M_n)_{n \in \mathbb{Z}}$ of non-negative integers. By the definition, it is easy to see that $h(M) = h(N)$ if and only if they have the same Hilbert series. Moreover, we also have that $h(M) = h(N)$ if and only if $h(M^*) = h(N^*)$ where $(-)^* = ^*\text{Hom}_R(-, \omega_R)$. See [7, Theorem 4.4.5.].

Remark 2.1. If M and N give the same class in the Grothendieck group, *i.e.* $[M] = [N]$ as an element of $K_0(\text{mod}^{\mathbb{Z}}(R))$ then $h(M) = h(N)$. However the converse does not hold in general. Let $R = k[x, y]/(x^2 - y^2)$ with $\deg x = \deg y = 1$ and set $M = R/(x + y)$ and $N = R/(x - y)$. Then $h(M) = h(N)$ and $[M] \neq [N]$ in $K_0(\text{mod}^{\mathbb{Z}}(R))$. In fact, let \mathfrak{p} be an ideal $(x + y)R$. Then $\text{rank}_{R_{(\mathfrak{p})}} M_{(\mathfrak{p})} = 1$ and $\text{rank}_{R_{(\mathfrak{p})}} N_{(\mathfrak{p})} = 0$. Note that $[M] = [N]$ on $K_0(\text{mod}^{\mathbb{Z}}(R_{(\mathfrak{p})}))$ yields that $[M_{(\mathfrak{p})}] = [N_{(\mathfrak{p})}]$ on $K_0(\text{mod}^{\mathbb{Z}}(R_{(\mathfrak{p})}))$. Thus $[M] = [N]$ can never happen.

Our motivation of the paper is to investigate the graded degenerations of graded Cohen-Macaulay modules in terms of some order relations. For the reason we consider the following relation on $\text{CM}^{\mathbb{Z}}(R)$ that is known as the hom order.

Definition 2.2. For $M, N \in \text{CM}^{\mathbb{Z}}(R)$ we define $M \leq_{\text{hom}} N$ if $[M, X] \leq [N, X]$ for each $X \in \text{CM}^{\mathbb{Z}}(R)$. Here $[M, X]$ is an abbreviation of $\dim_k \text{Hom}_R(M, X)$.

Remark 2.3. For $M, N \in \text{mod}^{\mathbb{Z}}(R)$, $[M, N]$ is finite, and thus we can consider the above relation. As a consequence in [5], if R is of dimension 0, 1 or 2, \leq_{hom} is a partial order on $\text{CM}^{\mathbb{Z}}(R)$ since $\text{CM}^{\mathbb{Z}}(R)$ is closed under kernels in such cases.

Lemma 2.4. *Let R be a graded Gorenstein ring and let M and N be indecomposable graded Cohen-Macaulay R -modules. Suppose that $h(M) = h(N)$. For $Y \in \text{mod}^{\mathbb{Z}}(R)$ which is of finite projective dimension, we have $[M, Y] = [N, Y]$.*

Proof. For a graded R -module Y which is of finite projective dimension, $\Omega^i Y$ is also of finite projective dimension. Since R is Gorenstein, we have $\text{Ext}_R^1(M, \Omega^i Y) = 0$ for all graded Cohen-Macaulay R -modules M . Thus, take a graded free resolution of Y and apply $\text{Hom}_R(M, -)$ to the resolution, we get an exact sequence

$$0 \rightarrow \text{Hom}_R(M, F_l) \rightarrow \text{Hom}_R(M, F_{l-1}) \rightarrow \cdots \rightarrow \text{Hom}_R(M, F_0) \rightarrow \text{Hom}_R(M, Y) \rightarrow 0.$$

Hence

$$[M, Y] = \sum_{i=0}^l (-1)^i [M, F_i].$$

We also have

$$[N, Y] = \sum_{i=0}^l (-1)^i [N, F_i].$$

Since R is Gorenstein, $\omega_R = R(l)$ for some $l \in \mathbb{Z}$. Then we have an equality $[M, F] = [N, F]$ for each free module. Therefore we have

$$[M, Y] = \sum_{i=0}^l (-1)^i [M, F_i] = \sum_{i=0}^l (-1)^i [N, F_i] = [N, Y].$$

□

In the paper we use the theory of Auslander-Reiten (abbr. AR) sequences of graded Cohen-Macaulay modules. For the detail, we recommend the reader to look at [9, 3, 4] and [13, Chapter 15.]. We denote by $\text{CM}_0^{\mathbb{Z}}(R)$ the full subcategory of $\text{CM}^{\mathbb{Z}}(R)$ consisting of $M \in \text{CM}^{\mathbb{Z}}(R)$ such that $M_{(\mathfrak{p})}$ is $R_{(\mathfrak{p})}$ -free for any graded prime ideal $\mathfrak{p} \neq \mathfrak{m}$.

Theorem 2.5. [3, 4, 13, 9] *Let (R, \mathfrak{m}) be a Noetherian \mathbb{Z} -graded Gorenstein $*$ -local ring. Then $\text{CM}_0^{\mathbb{Z}}(R)$ admits AR sequences.*

Definition 2.6. We say that (R, \mathfrak{m}) is a graded isolated singularity if each graded localization $R_{(\mathfrak{p})}$ is regular for each graded prime ideal \mathfrak{p} with $\mathfrak{p} \neq \mathfrak{m}$.

It is easy to see that $\text{CM}^{\mathbb{Z}}(R) = \text{CM}_0^{\mathbb{Z}}(R)$ if R is a graded isolated singularity, so that $\text{CM}^{\mathbb{Z}}(R)$ admits AR sequences. We denote by $\mu(M, Z)$ the multiplicity of Z as a direct summand of M .

Theorem 2.7. *Let R be a graded Gorenstein ring with R_0 is an algebraically closed field and let M and N graded Cohen-Macaulay R -modules. Assume that R is a graded isolated singularity. Then $[M, X] = [N, X]$ for each $X \in \text{CM}^{\mathbb{Z}}(R)$ if and only if $M \cong N$. Particularly, \leq_{hom} is a partial order on $\text{CM}^{\mathbb{Z}}(R)$.*

Proof. We decompose M as $M = \bigoplus M_i^{\mu(M, M_i)}$ where M_i are indecomposable graded Cohen-Macaulay R -modules. If M_i is not free, we can take the AR sequence ending in M_i

$$0 \rightarrow \tau M_i \rightarrow E_i \rightarrow M_i \rightarrow 0,$$

where τM_i is an AR translation of M_i . Apply $\text{Hom}_R(M, -)$ and $\text{Hom}_R(N, -)$ to the sequence, since k is an algebraically closed field, we have

$$0 \rightarrow \text{Hom}_R(M, \tau M_i) \rightarrow \text{Hom}_R(M, E_i) \rightarrow \text{Hom}_R(M, M_i) \rightarrow k^{\mu(M, M_i)} \rightarrow 0$$

and

$$0 \rightarrow \text{Hom}_R(N, \tau M_i) \rightarrow \text{Hom}_R(N, E_i) \rightarrow \text{Hom}_R(N, M_i) \rightarrow k^{\mu(N, M_i)} \rightarrow 0.$$

Counting the dimensions of terms, we conclude that $\mu(M, M_i) = \mu(N, M_i)$.

If M_i is free, we may assume that $M_i = R$. Let \mathfrak{m} be a *maximal ideal of R . We consider the Cohen-Macaulay approximation (see [2]) of \mathfrak{m}

$$0 \rightarrow Y \rightarrow X \rightarrow \mathfrak{m} \rightarrow 0.$$

We note that X is a graded Cohen-Macaulay R -module and Y is of finite injective dimension. We also note that Y is of finite projective dimension since R is Gorenstein. Let f be a composition map of the approximation $X \rightarrow \mathfrak{m}$ and a natural inclusion $\mathfrak{m} \rightarrow R$. Then we get the following commutative diagram.

$$\begin{array}{ccccccc} 0 & \longrightarrow & K & \longrightarrow & X & \xrightarrow{f} & R \\ & & \uparrow g & & \uparrow = & & \uparrow \subseteq \\ 0 & \longrightarrow & Y & \longrightarrow & X & \longrightarrow & \mathfrak{m} \longrightarrow 0. \end{array}$$

By a diagram chasing we see that g is surjective, so that K is of finite projective dimension. Now we obtain an exact sequence

$$0 \rightarrow \text{Hom}_R(M, K) \rightarrow \text{Hom}_R(M, X) \rightarrow \text{Hom}_R(M, R) \rightarrow k^{\mu(M, R)} \rightarrow 0.$$

According to Lemma 2.4, $[M, K] = [N, K]$. Thus

$$[M, R] + [M, K] - [M, X] = [N, R] + [N, K] - [N, X].$$

Hence $\mu(M, R) = \mu(N, R)$. Consequently $M \cong N$. □

3. GRADED DEGENERATIONS OF GRADED COHEN-MACAULAY MODULES

We define a notion of degenerations for graded modules.

Definition 3.1. Let R be a Noetherian \mathbb{N} -graded ring where $R_0 = k$ is a field and let $V = k[[t]]$ with a trivial gradation and K be the localization by t , namely $K = V_t = k(t)$. For finitely generated graded R -modules M and N , we say that M gradually degenerates to N or N is a graded degeneration of M if there is a finitely generated graded $R \otimes_k V$ -module Q which satisfies the following conditions:

- (1) Q is flat as a V -module.
- (2) $Q \otimes_V V/tV \cong N$ as a graded R -module.
- (3) $Q \otimes_V K \cong M \otimes_k K$ as a graded $R \otimes_k K$ -module.

In [15], Yoshino gives a necessary and sufficient condition for degenerations of (non-graded) modules. One can also show its graded version in a similar way. See also [12, 19].

Theorem 3.2. [15, Theorem 2.2] *The following conditions are equivalent for finitely generated graded R -modules M and N .*

- (1) M gradually degenerates to N .
- (2) *There is a short exact sequence of finitely generated graded R -modules*

$$0 \longrightarrow Z \longrightarrow M \oplus Z \longrightarrow N \longrightarrow 0.$$

Remark 3.3. (1) As Yoshino has shown in [15], the endomorphism of Z in the sequence of Theorem 3.2 is nilpotent. Note that we do not need the nilpotency assumption here. Actually, since $\text{End}_R(Z)$ is Artinian, by using Fitting theorem, we can describe the endomorphism as a direct sum of an isomorphism and a nilpotent morphism. See also [15, Remark 2.3.].

- (2) Assume that M and N are graded Cohen-Macaulay modules. Then we can show that Z is also a graded Cohen-Macaulay R -module. See [15, Remark 4.3.].
- (3) Assume that there is an exact sequence of finitely generated graded R -modules

$$0 \longrightarrow L \longrightarrow M \longrightarrow N \longrightarrow 0.$$

Then M gradually degenerates to $L \oplus N$. See [15, Remark 2.5] for instance.

- (4) Let M and N be finitely generated graded R -modules and suppose that M gradually degenerates to N . Then the modules M and N give the same class in the Grothendieck group.

We can also prove in a similar way to the proof of [17, Theorem 2.1.] that, for $L, M, N \in \text{mod}^{\mathbb{Z}}(R)$, if L gradually degenerates to M and if M gradually degenerates to N then L gradually degenerates to N . And one can show that if L gradually degenerates to M then $L \leq_{\text{hom}} M$. Consideration into this fact we define partial orders as follows.

Definition 3.4. For finitely generated graded R -modules M, N , we define the relation $M \leq_{deg} N$, which is called the degeneration order, if M gradually degenerates to N . We also define the relation $M \leq_{ext} N$ if there are modules M_i, N'_i, N''_i and short exact sequences $0 \rightarrow N'_i \rightarrow M_i \rightarrow N''_i \rightarrow 0$ in $\text{CM}^{\mathbb{Z}}(R)$ so that $M = M_1, M_{i+1} = N'_i \oplus N''_i, 1 \leq i \leq s$ and $N = M_{s+1}$ for some s .

For $M \in \text{CM}^{\mathbb{Z}}(R)$, we take a first syzygy module of M^*

$$0 \rightarrow \Omega^1 M^* \rightarrow F \rightarrow M^* \rightarrow 0.$$

Applying ${}^*\text{Hom}_R(-, \omega_R)$ to the sequence, we have

$$0 \rightarrow M^{**} \cong M \rightarrow F^* \rightarrow (\Omega^1 M^*)^* \rightarrow 0.$$

Then we denote $(\Omega^1 M^*)^*$ by $\Omega^{-1} M$.

Lemma 3.5. *Let R be a graded isolated singularity with ω_R and let M and N be graded Cohen-Macaulay R -modules. Assume that $h(M) = h(N)$ and $M \leq_{hom} N$. Then, for each graded Cohen-Macaulay R -module X , there exists an integer $l_X \gg 0$ such that $[M, X(\pm l_X)] = [N, X(\pm l_X)]$.*

Proof. For each $X \in \text{CM}^{\mathbb{Z}}(R)$, we can take an exact sequence as above

$$0 \rightarrow X \rightarrow E \rightarrow \Omega^{-1} X \rightarrow 0.$$

Note that E is a direct sum of $\omega_R(n)$ for some integers n . Applying ${}^*\text{Hom}_R(M, -)$ to the sequence, we have

$$0 \rightarrow {}^*\text{Hom}_R(M, X) \rightarrow {}^*\text{Hom}_R(M, E) \rightarrow {}^*\text{Hom}_R(M, \Omega^{-1} X) \rightarrow {}^*\text{Ext}_R(M, X) \rightarrow 0.$$

Since R is a graded isolated singularity, $\dim {}^*\text{Ext}_R(M, X)$ is finite. Thus ${}^*\text{Ext}_R(M, X)_{\pm l_1} = 0$ for sufficiently large $l_1 \gg 0$. Similarly there also exists an integer $l_2 \gg 0$ so that ${}^*\text{Ext}_R(N, X)_{\pm l_2} = 0$.

Set $l = \max\{l_1, l_2\}$. Then we have

$$0 \rightarrow {}^*\text{Hom}_R(M, X)_{\pm l} \rightarrow {}^*\text{Hom}_R(M, E)_{\pm l} \rightarrow {}^*\text{Hom}_R(M, \Omega^{-1} X)_{\pm l} \rightarrow 0.$$

Therefore we obtain the equation

$$[M, X(\pm l)] = [M, E(\pm l)] - [M, \Omega^{-1} X(\pm l)].$$

We also have

$$[N, X(\pm l)] = [N, E(\pm l)] - [N, \Omega^{-1} X(\pm l)].$$

Suppose that $[M, X(\pm l)] < [N, X(\pm l)]$. Then the following inequality holds.

$$[M, E(\pm l)] - [M, \Omega^{-1} X(\pm l)] < [N, E(\pm l)] - [N, \Omega^{-1} X(\pm l)].$$

Since $h(M) = h(N)$, $[M, E(\pm l)] = [N, E(\pm l)]$. Hence we see that $[M, \Omega^{-1} X(\pm l)] > [N, \Omega^{-1} X(\pm l)]$. This is a contradiction since $M \leq_{hom} N$. Therefore we have some integer l such that $[M, X(\pm l)] = [N, X(\pm l)]$. \square

We say that the category $\text{CM}^{\mathbb{Z}}(R)$ is of graded finite representation type if there are only a finite number of isomorphism classes of indecomposable graded Cohen-Macaulay modules up to shift. We note that if $\text{CM}^{\mathbb{Z}}(R)$ is of finite representation type, then R is a graded isolated singularity. See [13, Chapter 15.] for the detail.

As an immediate consequence of Lemma 3.5, we have the following.

Corollary 3.6. *Let R be of finite representation type and let M and N be graded Cohen-Macaulay R -modules. Assume that $h(M) = h(N)$ and $M \leq_{\text{hom}} N$. Then there are only finitely many indecomposable graded Cohen-Macaulay R -modules X such that $[N, X] - [M, X] > 0$.*

For graded Cohen-Macaulay R -modules M and N , we consider the following set of all the isomorphism classes of indecomposable graded Cohen-Macaulay modules

$$\mathcal{F}_{M,N} = \{X \mid [N, X] - [M, X] > 0\} / \cong.$$

Note from Corollary 3.6 that \mathcal{F} is a finite set if $h(M) = h(N)$ and $M \leq_{\text{hom}} N$. We also note that $\omega_R \notin \mathcal{F}$ in the case.

Proposition 3.7. [12] *Let R be a graded Gorenstein ring which is of graded finite representation type and let M and N be graded Cohen-Macaulay R -modules. Assume that $h(M) = h(N)$ and $M \leq_{\text{hom}} N$. Then there exists some graded Cohen-Macaulay R -module L such that $M \oplus L$ degenerates to $N \oplus L$.*

Proof. Although a proof of the proposition is given in [12], we refer the argument of the proof in the present paper. For this reason we briefly recall the proof of the proposition.

Since R is a graded isolated singularity, $\text{CM}^{\mathbb{Z}}(R)$ admits AR sequences. For each $X \in \mathcal{F}_{M,N}$, we can take an AR sequence starting from X .

$$\Sigma_X : 0 \rightarrow X \rightarrow E_X \rightarrow \tau^{-1}X \rightarrow 0.$$

Now we consider a sequence which is a direct sum of $[N, X] - [M, X]$ copies of Σ_X where X runs through all modules in $\mathcal{F}_{M,N}$. Namely

$$\bigoplus_{X \in \mathcal{F}_{M,N}} \Sigma_X^{[N, X] - [M, X]} = 0 \rightarrow U \rightarrow V \rightarrow W \rightarrow 0.$$

For any indecomposable $Z \in \text{CM}^{\mathbb{Z}}(R)$, we obtain

$$0 \rightarrow \text{Hom}_R(W, Z) \rightarrow \text{Hom}_R(V, Z) \rightarrow \text{Hom}_R(U, Z) \rightarrow k^{[N, Z] - [M, Z]} \rightarrow 0.$$

This implies that the equality

$$[U, Z] + [W, Z] - [V, Z] = [N, Z] - [M, Z],$$

thus

$$[U, Z] + [W, Z] + [M, Z] = [N, Z] + [V, Z]$$

holds for all $Z \in \text{CM}^{\mathbb{Z}}(R)$. Hence this yields that

$$M \oplus U \oplus W \cong N \oplus V.$$

Since V degenerates to $U \oplus W$, therefore $M \oplus V$ degenerates to $M \oplus U \oplus W \cong N \oplus V$. \square

Now we focus on the case that a graded Gorenstein ring is of graded finite representation type and representation directed. We say that a graded Cohen-Macaulay ring R is representation directed if the AR quiver of $\text{CM}_0^{\mathbb{Z}}(R)$ has no oriented cyclic paths. Bongartz [6] has studied such a case over finite dimensional k -algebras. In our graded settings, the similar results hold. Actually we shall prove the following.

Theorem 3.8. *Let R be a graded Gorenstein ring which is of graded finite representation type and representation directed. Then the following conditions are equivalent for M and $N \in \text{CM}^{\mathbb{Z}}(R)$.*

- (1) $h(M) = h(N)$ and $M \leq_{\text{hom}} N$.
- (2) $M \leq_{\text{deg}} N$.
- (3) $M \leq_{\text{ext}} N$.

To prove the theorem, we modify the arguments in [6].

Lemma 3.9 (Cancellation property). *Let M , N and X be finitely generated graded R -modules.*

- (1) *Assume that $[X, M] = [X, N]$. If $M \oplus X$ gradually degenerates to $N \oplus X$, M gradually degenerates to N .*
- (2) *Assume that R is Gorenstein and M and N graded Cohen-Macaulay R -modules. If M gradually degenerates to $N \oplus F$ for some graded free R -module F then M/F gradually degenerates to N .*

Proof. (1) Since $M \oplus X$ gradually degenerates to $N \oplus X$, there exist an exact sequence

$$0 \rightarrow W \rightarrow M \oplus X \oplus W \rightarrow N \oplus X \rightarrow 0.$$

We construct a pushout diagram.

$$\begin{array}{ccccccc}
 & & & 0 & & 0 & \\
 & & & \uparrow & & \uparrow & \\
 & & & X & \xlongequal{\quad} & X & \\
 & & & \uparrow & & \uparrow & \\
 0 & \longrightarrow & W & \longrightarrow & M \oplus X \oplus W & \longrightarrow & N \oplus X \longrightarrow 0 \\
 & & \parallel & & \uparrow & & \uparrow \\
 0 & \longrightarrow & W & \longrightarrow & E & \longrightarrow & N \longrightarrow 0 \\
 & & & & \uparrow & & \uparrow \\
 & & & & 0 & & 0.
 \end{array}$$

For the middle column sequence,

$$[X, X] + [X, E] - [X, M \oplus X \oplus W] = [X, E] - [X, M] - [X, W] \geq 0.$$

On the other hand, for the bottom row sequence, since $[X, M] = [X, N]$,

$$[X, N] + [X, W] - [X, E] = [X, M] + [X, W] - [X, E] \geq 0.$$

Thus we have

$$[X, X] + [X, E] - [X, M \oplus X \oplus W] = 0.$$

This implies that the middle column sequence splits, so that $X \oplus E \cong M \oplus X \oplus W$. Therefore $E \cong M \oplus W$ and we get

$$0 \rightarrow W \rightarrow M \oplus W \rightarrow N \rightarrow 0.$$

Namely M gradually degenerates to N .

(2) Since M gradually degenerates to $N \oplus F$, we have an exact sequence

$$0 \rightarrow Z \rightarrow M \oplus Z \rightarrow N \oplus F \rightarrow 0.$$

Suppose that Z contains a graded free module G as a direct summand. We construct a pushout diagram

$$\begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \uparrow & & \uparrow & & \\ 0 & \longrightarrow & Z/G & \longrightarrow & E & \longrightarrow & N \oplus F \longrightarrow 0 \\ & & \uparrow & & \uparrow & & \parallel \\ 0 & \longrightarrow & Z & \longrightarrow & M \oplus Z & \longrightarrow & N \oplus F \longrightarrow 0 \\ & & \uparrow & & \uparrow & & \\ & & G & \xlongequal{\quad} & G & & \\ & & \uparrow & & \uparrow & & \\ & & 0 & & 0 & & \end{array}$$

The left column sequence is a split sequence induced by the decomposition $Z \cong Z/G \oplus G$. Note that E is also a graded Cohen-Macaulay module. Since R is Gorenstein, the middle column sequence is also split. Hence

$$E \oplus G \cong M \oplus Z \cong M \oplus Z/G \oplus G.$$

This yields that $E \cong M \oplus Z/G$. Hence we may assume that Z has no graded free modules as direct summands.

Consider a composition of the surjection $M \oplus Z \rightarrow N \oplus F$ and the projection $N \oplus F \rightarrow F$. Then the composition mapping is split, so that M contains F as a direct summand. Hence $M \cong M/F \oplus F$ gradually degenerates to $N \oplus F$. Since $[F, M/F \oplus F] = [F, M] = [F, N \oplus F]$, by (1), we conclude that M/F gradually degenerates to N . \square

For indecomposable graded Cohen-Macaulay modules M and N , we write $X \preceq Y$ if $X \cong Y$ or if there exists a finite path from X to Y in the AR quiver of $\text{CM}_0^{\mathbb{Z}}(R)$.

Lemma 3.10. *Let R be a graded Gorenstein ring which is of graded finite representation type and representation directed and let M and $N \in \text{CM}^{\mathbb{Z}}(R)$. Assume that $h(M) = h(N)$, $M \leq_{\text{hom}} N$ and M and N have no common direct summands. Let $X \in \text{CM}^{\mathbb{Z}}(R)$ be an indecomposable module such that \preceq -minimal with the property $[N, X] - [M, X] > 0$ and E be a middle term of an AR sequence starting from X . Then $[E, N] = [E, M]$.*

Proof. As in the proof of Proposition 3.7, we can construct the sequence

$$0 \rightarrow U \rightarrow V \rightarrow W \rightarrow 0$$

in $\text{CM}^{\mathbb{Z}}(R)$ such that $U \oplus W \oplus M \cong V \oplus N$ via taking a direct sum of AR sequences starting from modules in $\mathcal{F}_{M,N}$. This isomorphism implies that $[Z, U] + [Z, W] - [Z, V] = [Z, N] - [Z, M]$ for each $Z \in \text{CM}^{\mathbb{Z}}(R)$. Thus it is enough to show that the equality $[E, U] + [E, W] - [E, V] = 0$ holds. If there is a $Y \in \mathcal{F}_{M,N}$ such that, for the AR sequence $0 \rightarrow Y \rightarrow G \rightarrow \tau^{-1}Y \rightarrow 0$,

$$[E, Y] + [E, \tau^{-1}Y] - [E, G] > 0.$$

If $Y \cong X$, then one can show that $\tau^{-1}X$ is a direct summand of E , so that there is a cyclic path. This is a contradiction since R is representation directed. Thus Y is not isomorphic to X . The inequality also show that $\tau^{-1}Y$ is a direct summand of E . Thus there is an irreducible map from X to $\tau^{-1}Y$. Hence X is a direct summand of G , so that $Y \preceq X$. This is a contradiction. Consequently, for each $Y \in \mathcal{F}_{M,N}$, $[E, Y] + [E, \tau^{-1}Y] - [E, G] = 0$, therefore $[E, N] = [E, M]$. \square

Proof of Theorem 3.8. The implication (3) \Rightarrow (2) \Rightarrow (1) is trivial. Note that $\text{Ext}_R(X, X) = 0$ for an indecomposable $X \in \text{CM}^{\mathbb{Z}}(R)$ since R is representation directed. Thus the implication (2) \Rightarrow (3) can be shown by the same argument in [18, 3.5.]. Now we shall show (1) \Rightarrow (2). Set $V = \bigoplus_{W \in \mathcal{F}_{M,N}} W$. Since $[N, X] - [M, X] = 0$ for any $X \notin \mathcal{F}_{M,N}$, we can show the implication by induction on $d = [N, V] - [M, V]$. If $d = 0$, $[N, X] = [M, X]$ for each $X \in \text{CM}^{\mathbb{Z}}(R)$. Thus $M \cong N$. Hence assume that $d > 0$ and M and N have no summand in common in the inductive step. We take $X \in \text{CM}^{\mathbb{Z}}(R)$ in Lemma 3.10 and let E be a middle term of the AR sequence starting from X . By virtue of Lemma 3.10 and Lemma 3.9 (1) it is enough to show that $E \oplus M \leq_{\text{deg}} E \oplus N$. Now we have $E \oplus M \leq_{\text{ext}} X \oplus \tau^{-1}X \oplus M$. By the property of AR sequence, for each indecomposable Y ,

$$[X, Y] + [\tau^{-1}X, Y] = [E, Y] + \delta_{X,Y}$$

where $\delta_{X,Y} = 1$ if $X \cong Y$ and otherwise 0. Thus $X \oplus \tau^{-1}X \oplus M \leq_{\text{hom}} E \oplus N$. We should remark that $\mathcal{F}_{X \oplus \tau^{-1}X \oplus M, E \oplus N}$ is contained in $\mathcal{F}_{M,N}$ since $X \in \mathcal{F}_{M,N}$.

Then

$$\begin{aligned} & [E \oplus N, V] - [X \oplus \tau^{-1}X \oplus M, V] \\ &= [N, V] - [M, V] + [E, V] - [X \oplus \tau^{-1}X, V] \\ &= d - 1 \end{aligned}$$

By the induction hypothesis,

$$E \oplus M \leq_{\text{ext}} X \oplus \tau^{-1}X \oplus M \leq_{\text{deg}} E \oplus N$$

so that $E \oplus M \leq_{\text{deg}} E \oplus N$. \square

Remark 3.11. The implication $M \leq_{\text{ext}} N \Rightarrow M \leq_{\text{deg}} N$ does not hold in general. Let $R = k[x, y]/(x^2)$ with $\deg x = \deg y = 1$. Then $R(-1)$ gradually degenerates to $(x, y^2)R$. In fact we have an exact sequence

$$0 \longrightarrow R/(x)(-2) \xrightarrow{\begin{pmatrix} x \\ 0 \end{pmatrix}} R(-1) \oplus R/(x)(-2) \xrightarrow{\begin{pmatrix} x & y^2 \end{pmatrix}} (x, y^2)R \longrightarrow 0.$$

Since $(x, y^2)R$ is an indecomposable graded Cohen-Macaulay module which is not isomorphic to $R(-1)$, $R(-1) \leq_{\text{ext}} (x, y^2)R$ can never happen. See also [8, Remark 2.5.].

Proposition 3.12. *Let R be a graded Gorenstein ring which is of graded finite representation type and let M and N be graded Cohen-Macaulay R -modules. Assume that $h(M) = h(N)$ and $M \leq_{\text{hom}} N$. Then for each indecomposable non-free graded Cohen-Macaulay R -module X we have the following equality.*

$$[N, X] - [M, X] = [\tau^{-1}X, N] - [\tau^{-1}X, M].$$

Proof. Under the assumption, as in the proof of Proposition 3.7, we can construct an exact sequence

$$0 \rightarrow U \rightarrow V \rightarrow W \rightarrow 0$$

in $\text{CM}^{\mathbb{Z}}(R)$ such that $[U, X] + [W, X] - [V, X] = [N, X] - [M, X]$. It is enough to show that the equality $[U \oplus W, X] - [V, X] = [\tau^{-1}X, U \oplus W] - [\tau^{-1}X, V]$ holds for each indecomposable X . Moreover the sequence is a direct sum of AR sequences, we may assume that $0 \rightarrow U \rightarrow V \rightarrow W \rightarrow 0$ is an AR sequence starting from U . Let $X \in \text{CM}^{\mathbb{Z}}(R)$ be indecomposable and non projective. By the property of an AR sequence, we have

$$[U \oplus W, X] - [V, X] = \delta_{U, X} \quad [\tau^{-1}X, U \oplus W] - [\tau^{-1}X, V] = \delta_{W, \tau^{-1}X}.$$

Since $W \cong \tau^{-1}U$, we can get the equality. \square

4. REMARKS ON STABLE DEGENERATIONS OF GRADED COHEN-MACAULAY MODULES

In the rest of the paper we consider the stable analogue of degenerations of graded Cohen-Macaulay modules.

Let R be a graded Gorenstein ring where $R_0 = k$ is a field and let $V = k[[t]]$ with a trivial gradation and $K = k(t)$. Note that $R \otimes_k V$ and $R \otimes_k K$ are

graded Gorenstein rings as well. Then $\underline{\mathbf{CM}}^{\mathbb{Z}}(R \otimes_k V)$ and $\underline{\mathbf{CM}}^{\mathbb{Z}}(R \otimes_k K)$ are triangulated categories. We denote by $\mathbf{L} : \underline{\mathbf{CM}}^{\mathbb{Z}}(R \otimes_k V) \rightarrow \underline{\mathbf{CM}}^{\mathbb{Z}}(R \otimes_k K)$ (resp. $\mathcal{R} : \underline{\mathbf{CM}}^{\mathbb{Z}}(R \otimes_k V) \rightarrow \underline{\mathbf{CM}}^{\mathbb{Z}}(R)$) the triangle functor defined by the localization by t (resp. taking $-\otimes_V V/tV$). See also [16, Definition 4.1].

Definition 4.1. Let $\underline{M}, \underline{N} \in \underline{\mathbf{CM}}^{\mathbb{Z}}(R)$. We say that \underline{M} stably degenerates to \underline{N} if there exists a graded Cohen-Macaulay module $\underline{Q} \in \underline{\mathbf{CM}}^{\mathbb{Z}}(R \otimes_k V)$ such that $\mathbf{L}(\underline{Q}) \cong \underline{M} \otimes_k K$ in $\underline{\mathbf{CM}}^{\mathbb{Z}}(R \otimes_k K)$ and $\mathcal{R}(\underline{Q}) \cong \underline{N}$ in $\underline{\mathbf{CM}}^{\mathbb{Z}}(R)$.

One can show the following characterization of stable degenerations similarly to the proof of [16, Theorem 5.1].

Theorem 4.2. *Let R be a graded Gorenstein ring where $R_0 = k$ is a field. The following conditions are equivalent for graded Cohen-Macaulay R -modules M and N .*

- (1) $F \oplus M$ degenerates to N for some graded free R -module F .
- (2) There is a triangle in $\underline{\mathbf{CM}}^{\mathbb{Z}}(R)$

$$\underline{Z} \longrightarrow \underline{M} \oplus \underline{Z} \longrightarrow \underline{N} \longrightarrow \underline{Z}[1].$$

- (3) \underline{M} stably degenerates to \underline{N} .

Proof. We should note that the implication (3) \Rightarrow (1). In our setting, $R \otimes_k V$ and $R \otimes_k K$ are * local. Then graded projective $R \otimes_k V$ (resp. $R \otimes_k K$) -modules are graded free $R \otimes_k V$ (resp. $R \otimes_k K$) -modules. Hence we can show the implication as in the Artinian case of the proof of [16, Theorem 5.1]. \square

Remark 4.3. A theory of degenerations for derived categories has been studied in [10]. They have shown that, for complexes M, N in the bounded derived category of a finite dimensional algebra, M degenerates to N if and only if there exists a triangle of the form which appears in the above theorem. Let R be a graded Gorenstein ring with $R_0 = k$ is an algebraically closed field. As shown in [1, 11, 9], suppose that R has a simple singularity then there exists a Dynkin quiver Q such that we have a triangle equivalence

$$\underline{\mathbf{CM}}^{\mathbb{Z}}(R) \cong \mathbf{D}^b(kQ)$$

where $\mathbf{D}^b(kQ)$ is a bounded derived category of the category of finitely generated left modules over a path algebra kQ . By virtue of Theorem 4.2, we can describe the degenerations for $\mathbf{D}^b(kQ)$ in terms of the graded degenerations for $\underline{\mathbf{CM}}^{\mathbb{Z}}(R)$. Since the graded ring R is of graded finite representation type and representation directed, we have already seen them in Theorem 3.8.

ACKNOWLEDGMENTS

The author express his deepest gratitude to Tokuji Araya and Yuji Yoshino for valuable discussions and helpful comments.

REFERENCES

- [1] T. ARAYA, *Exceptional sequences over graded Cohen-Macaulay rings*. Math. J. Okayama Univ. **41** (1999), 81–102 (2001).
- [2] M. AUSLANDER and R.-O. BUCHWEITZ, *The homological theory of maximal Cohen-Macaulay approximations*. Mem. Soc. Math. France (N.S.) No. 38 (1989), 5–37.
- [3] M. AUSLANDER and I. REITEN, *Almost split sequences for \mathbb{Z} -graded rings*. Singularities, representation of algebras, and vector bundles (Lambrecht, 1985), 232–243, Lecture Notes in Math., 1273, Springer, Berlin, 1987.
- [4] M. AUSLANDER and I. REITEN, *Almost split sequences for abelian group graded rings*. J. Algebra **114** (1988), no. 1, 29–39.
- [5] K. BONGARTZ, *A generalization of a theorem of M. Auslander*. Bull. London Math. Soc. **21** (1989), no. 3, 255–256.
- [6] K. BONGARTZ, *On degenerations and extensions of finite-dimensional modules*. Adv. Math. **121** (1996), 245–287.
- [7] W. BRUNS and J. HERZOG, *Cohen-Macaulay Rings*, Cambridge Studies in Advanced Mathematics, 39. Cambridge University Press, Cambridge, 1993. xii+403 pp. Revised edition, 1998.
- [8] N. HIRAMATSU and Y. YOSHINO, *Examples of degenerations of Cohen-Macaulay modules*, to appear in Proc. Amer. Math. Soc., arXiv1012.5346.
- [9] O. IYAMA and R. TAKAHASHI, *Tilting and cluster tilting for quotient singularities*, to appear in Math. Ann., arXiv:1012.5954.
- [10] B. JENSEN, X. SU and A. ZIMMERMANN, *Degenerations for derived categories*. J. Pure Appl. Algebra **198** (2005), no. 1-3, 281–295.
- [11] H. KAJIURA, K. SAITO and A. TAKAHASHI, *Matrix factorization and representations of quivers. II. Type ADE case*. Adv. Math. **211** (2007), no. 1, 327–362.
- [12] C. RIEDTMANN, *Degenerations for representations of quivers with relations*. Ann. Scient. École Norm. Sup. 4^e série **19** (1986), 275–301.
- [13] Y. YOSHINO, *Cohen-Macaulay Modules over Cohen-Macaulay Rings*, London Mathematical Society Lecture Note Series **146**. Cambridge University Press, Cambridge, 1990. viii+177 pp.
- [14] Y. YOSHINO, *On degenerations of Cohen-Macaulay modules*. J. Algebra **248** (2002), 272–290.
- [15] Y. YOSHINO, *On degenerations of modules*. J. Algebra **278** (2004), 217–226.
- [16] Y. YOSHINO, *Stable degenerations of Cohen-Macaulay modules*, J. Algebra **332** (2011), 500–521.
- [17] G. ZWARA, *A degeneration-like order for modules*. Arch. Math. **71** (1998), 437–444.
- [18] G. ZWARA, *Degenerations for modules over representation-finite algebras*. Proc. Amer. Math. Soc. **127** (1999), 1313–1322.
- [19] G. ZWARA, *Degenerations of finite-dimensional modules are given by extensions*. Compositio Math. **121** (2000), 205–218.

DEPARTMENT OF GENERAL EDUCATION, KURE NATIONAL COLLEGE OF TECHNOLOGY,
2-2-11, AGAMINAMI, KURE HIROSHIMA, 737-8506 JAPAN

E-mail address: hiramatsu@kure-nct.ac.jp